

A case report of Hybrid verrucous carcinoma of the palate

Parya Emamverdyzade¹, Sona Rafieyan¹, Kuroush Taheritalesh², Ali Mortazavi^{2*}, Saeed Hajizadeh²

¹Department of Oral and Maxillofacial Pathology, Faculty of Dentistry, Tabriz University of Medical Sciences, Tabriz, Iran

²Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Tabriz University of Medical Sciences, Tabriz, Iran

*Corresponding Author:

Sona Rafieyan

Email: aliweb001@gmail.com

Abstract: Verrucous carcinoma (VC) a low-grade variant of squamous cell carcinoma (SCC) is the rarest of all oral cancers. A hybrid VC is a non-verrucous SCC that arises synchronously with the VC. The differential diagnosis of VC remains difficult and requires clinical and pathologic data confrontation. As the malignant behaviour of hybrid VC is confined to the non-VC component, careful examination of these tumors is needed. Here we report a case of 44 year old female with hybrid verrucous carcinoma. Clinical and histological features and treatment are discussed with the review of literature.

Keywords: verrucous carcinoma, hybrid verrucous carcinoma, palate

INTRODUCTION

It was first described in 1948 by Lauren V. Ackermann as a neoplasm of the oral mucous membrane, which is now also known as Verrucous Carcinoma of Ackermann or Ackermann's tumor. Tobacco chewing is a significant etiological factor for its development [1]. Other irritants to the oral mucosa such as betel nut chewing, poor dental hygiene and Human Papilloma Virus (HPV) infection have been implicated in the development of oral VC [2]. This uncommon lesion can be considered a disease of older age, typically occurring in the seventh-eighth decades, with a strong male predominance. In the head and neck area, VC most frequently involves the oral cavity, where it commonly seen in buccal mucosa and lip. It appears as a papillary nonulcerated gray-white or red mass with a very wide base of attachment [3]. Carcinomas composed of both verrucous carcinoma and conventional squamous cell carcinoma are referred to as hybrid verrucous squamous cell carcinoma (HCs) [4]. Although uncommon, it is estimated that about 10% of all verrucous carcinoma of the larynx and 20% of all verrucous carcinoma of the oral cavity are hybrid [4]. Careful examination is recommended because the malignant behaviour of hybrid VC is confined to the non-VC component [4].

CASE REPORT

A 44-year-old woman presented to the Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Tabriz University of Medical Sciences, Tabriz, Iran, with a chief complaint of an ulcer on the hard palate. The patient had a history of diabetes mellitus and renal failure. Clinical intraoral examination

revealed multiple sessile exophytic lesions with red dotted area measuring 15×9mm. It was painless and without bleeding. Paresthesia of inferior orbital nerve was observed. In radiographic studies no changes were seen. Computed tomography (CT) scan of this massive lesion did not show sinus involvement and bone erosion. The biopsy of lesion was done with differential diagnosis of verrucous carcinoma and sent to histopathological examination (Fig-1).



Fig-1: Multiple sessile exophytic lesions with red dotted area measuring 15×9mm

Histopathologic Findings

The histological appearance was described as well differentiated squamous tumor covered by a thick keratinized layer and papillary surface with broad and deep rete ridges. A typically inflammatory reaction in the stroma composed of lymphocytes, plasma cells and

histiocytes was observed in deep areas frank infiltration to the stroma, cytological atypia, individual cell keratinization were seen and the final diagnosis was hybrid verrucous carcinoma (Pictures-2&3).




Fig-2: In deep areas frank infiltration to the stroma is obvious (×400)




Fig-3: Tumor with broad and deep rete ridges (×200)

DISCUSSION

Verrucous carcinoma first described in 1948 by Lauren V. Ackerman is a distinct variant of differentiated SCC with low grade malignancy, slow growth and low metastatic potential [5]. The tumor representing 2–12% of all oral cancers mainly occurs in older men [6]. It is often associated with long-term use of smokeless tobacco. The VC most often arises in the upper aerodigestive tract also the oral cavity, particularly the buccal mucosa, gingivae and retromolar areas, remains the most common site of origin [7]. The etiology of VC remains unclear. The role of Human papillomavirus (HPV) infections in the etiology of verrucous lesion of the skin and genitalia is well documented, and HPV may also play an important role in the development of VC [2]. In clinical examination surface may be verrucous or show the conventional invasive pattern. To our knowledge it is important to take different incisions because in SCC with an exo-endophytic growth pattern often the invasion can be lacking in incisional biopsies, and it is not possible to exclude an underlying conventional carcinoma. Diagnosis from classical squamous cell

carcinoma is a frequent problem also for clinicians because of the extensive nature of the lesion mimicking an invasive cancer. In superficial biopsies without an obvious invasive growth, the benign microscopic appearance may, also, induce to an erroneous diagnosis of benign squamous proliferation [3]. In particular it is essential to rule out hybrid carcinoma including VC and conventional SCC. Hybrid carcinomas should be staged and managed as conventional SCC because of their metastatic potential, compare to classical VC and fortunately it shows excellent prognosis following complete surgical removal in the early stages [3]. Finally, it is essential that the pathologist alerts the clinician to the progressive nature of the lesion and because of the possibility of nodal metastasis complete excision or close followup and rebiopsy are suggested.

CONCLUSION

As the malignant behaviour of hybrid VC is confined to the non-VC component, careful examination of these tumors and adequate sampling is recommended.

REFERENCES

1. Kaushal, N., & Madan, N. (2011). Verrucous carcinoma of the oral cavity: case report. *The Internet Journal of Geriatrics and Gerontology*, 6(1).
2. Depprich, R. A., Handschel, J. G., Fritzemeier, C. U., Engers, R., & Kübler, N. R. (2006). Hybrid verrucous carcinoma of the oral cavity: A challenge for the clinician and the pathologist. *Oral Oncology Extra*, 42(2), 85-90.
3. Santoro, A., Pannone, G., Contaldo, M., Sanguedolce, F., Esposito, V., Serpico, R., ... & Bufo, P. (2010). A troubling diagnosis of verrucous squamous cell carcinoma ("the bad kind" of keratosis) and the need of clinical and pathological correlations: a review of the literature with a case report. *Journal of skin cancer*, 2011.
4. Barnes, L., Chiosea, S., Seethala, R.R. (2011). head and neck pathology. In: Barnes L, consultant pathology.
5. Steffen, C. (2004). The man behind the eponym: Lauren V. Ackerman and verrucous carcinoma of Ackerman. *The American Journal of Dermatopathology*, 26(4), 334-341.
6. Koch, B. B., Trask, D. K., Hoffman, H. T., Karnell, L. H., Robinson, R. A., Zhen, W., & Menck, H. R. (2001). National survey of head and neck verrucous carcinoma. *Cancer*, 92(1), 110-120.
7. Spiro, R. H. (1998). Verrucous carcinoma, then and now. *The American Journal of Surgery*, 176(5), 393-397.